

UNCLASSIFIED

AD 412108

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

AD NO. 412108

DDC FILE COPY

JPRS: 19,152

13 May 1963


OTS: 63-21005

LOWERING THE IMMUNOBIOLOGICAL RESISTANCE OF THE
ORGANISM AS A MANIFESTATION OF THE IRREVERSIBLE
COMPONENT OF RADIATION SICKNESS

by I. G. Akoyev, et al.

- USSR -

412108



U. S. DEPARTMENT OF COMMERCE
OFFICE OF TECHNICAL SERVICES
JOINT PUBLICATIONS RESEARCH SERVICE
Building T-30
Ohio Drive and Independence Avenue, S.W.
Washington 25, D. C.

Price: \$.50

FOREWORD

This publication was prepared under contract for the Joint Publications Research Service as a translation or foreign-language research service to the various federal government departments.

The contents of this material in no way represent the policies, views or attitudes of the U. S. Government or of the parties to any distribution arrangement.

PROCUREMENT OF JPRS REPORTS

All JPRS reports may be ordered from the Office of Technical Services. Reports published prior to 1 February 1963 can be provided, for the most part, only in photocopy (xerox). Those published after 1 February 1963 will be provided in printed form.

Details on special subscription arrangements for JPRS social science reports will be provided upon request.

No cumulative subject index or catalog of all JPRS reports has been compiled.

All JPRS reports are listed in the Monthly Catalog of U. S. Government Publications, available on subscription at \$4.50 per year (\$6.00 foreign), including an annual index, from the Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C.

All JPRS scientific and technical reports are cataloged and subject-indexed in Technical Translations, published ~~semi~~monthly by the Office of Technical Services, and also available on subscription (\$12.00 per year domestic, \$16.00 foreign) from the Superintendent of Documents. Semiannual indexes to Technical Translations are available at additional cost.

LOWERING THE IMMUNOBIOLOGICAL RESISTANCE OF THE
ORGANISM AS A MANIFESTATION OF THE IRREVERSIBLE
COMPONENT OF RADIATION SICKNESS

- USSR -

[Following is a translation of an article by I. G. Akoyev and M. A. Lagun in the Russian-language periodical Meditsinskaya Radiologiya (Medical Radiology), Vol VIII, No 2, Moscow, 1963, pages 47-50.]

According to hypotheses of I. F. Kovalyov and Blair -- Davidson (G. Davidson; V. N. Korogodin and G. G. Polikarpov; Blair, Storer, Harris, Furshner, Longham; Sparling, Strang, Worman), all affections of an organism caused by ionizing radiation can be divided into two types which are conditionally called reversible and irreversible. By this is meant not only truly irreversible damage (for example, genetic), but also that which is very slowly restored in comparison with reversible damage.

The reversible component of radiation sickness is associated in the first instance with damage to tissue systems which have a high level of physiological regeneration (hematogenic, epithelial, and others). Using the degree of reduction in the organism's immunity to re-irradiation as a criterion of the

amount of remaining radiation damage, it was established that the restoration period of radioresistance of an organism depends directly upon the species peculiarities and first and foremost upon the rate of metabolism processes. The restoration period of radioresistance (which is identified with the reparation period of all reversible injuries) for mice turned out to be 12 to 20 days (Kohn, Kallman, Rothermel, Woodard, Storer; Stearner, Tyler), and for man, according to the estimation by G. Davidson, this period stretches up to 200 days. According to our data, reparation of injuries in mice takes place in the course of 12 to 15 days. (Diagram 1 and Table 1).




Diagram 1. Dynamics of reparation of reversible radiation injuries of mice after irradiation, dose 400 rontgens. Criterion — displacement of average-lethal doses upon re-irradiation. (530 female mice, 2 to 3 months of age.)

Legend:

1. Residual radiation damage
2. Period of reduced radioresistance
3. Period of increased radioresistance
4. Days after irradiation

days. Upon double irradiation contamination occurred 50 days after the first irradiation. The animals were subjected to gamma rays Co^{60} , dosage rate 35r/minute. 150 white female mice ranging in age from two to three months were used in experiments.

The basic conditions of irradiation and results of the experiments are shown in Diagram 2 and in Table 2.




Diagram 2. Influence of conditions of preliminary irradiation on the life span of mice after contamination with toxin *Vas. perfringens* (contamination subcutaneous, dose 0.8 of experimental unit, 50 days after irradiation).

Probability of differences: Groups I and II, III, IV, V, VI -- $R = 0.99$; groups II and III, IV -- $R < 0.5$; groups V and VI -- $R < 0.5$; groups II + III + IV and V + VI $R = 0.99$.

- Legend: 1. Relative survival (%) 4. Time after toxin injection, minutes.
2. Average life span, minutes
3. Group of animals

① Group	② Dose of irradiation, r	③ Interval between irradiations	④ Equation of the curve	⑤ Average life span, minutes
I	0	—	$y = 187,5 - 178,5x$	$589 \pm 5,5$
II	750	—	$y = 130,0 - 125,2x$	384 ± 38
III	400 + 350	2	$y = 128,4 - 128,6x$	339 ± 13
IV	400 + 250	5	$y = 108,2 - 103,8x$	349 ± 13
V	400 + 500	8	$y = 101,6 - 119,9x$	$273 \pm 4,1$
VI	400 + 800	15	$y = 113,6 - 126,6x$	286 ± 14

Table 2

Legend:

1. Group
2. Irradiation dose
3. Interval between irradiations
4. Equation of the curve
5. Average life span, minutes

In all the groups 100% mortality of the preliminarily irradiated animals was observed. The average life span of the controlled animals was 589 ± 5 minutes, but of those exposed to single irradiation, dose 750 r — 384 ± 38 minutes. Consequently, even in the period of time after the restoration of the organism's radioresistance, the immunobiological state remains disrupted. Inasmuch as after irradiation there elapsed more than three times the time necessary for reparation of reversible radiation injuries, it is possible to suppose that reduced immunity of these animals to a bacterial toxin is linked with the irreversible component of radiation sickness. To substantiate this situation it was also necessary to show that the degree of lesion of immunobiological

resistance depends neither upon fractionation of the dose nor upon an increase in time intervals between irradiations. The following experiments were conducted to examine these problems.

In the second series of experiments the animals were subjected to double irradiation of the same total dosage — 750 r (400 + 350 r) — the time interval between exposures two and five days. In this short period of time about 60 and 75% of the reversible radiation injuries is repaired (cf. Diagram 1). The average life span of these mice after contamination by the toxin (339 ± 13 and 342 ± 13 minutes) essentially did not differ from the group exposed to single irradiation. Consequently, the degree of lesion of immunobiological resistance of the organism essentially does not depend upon the conditions of irradiation.

In the third series of experiments the influence of dosage and increased time interval between irradiations up to a period of time necessary for full reparation of reversible radiation alterations brought about by the first irradiation was examined. The animals of this series were exposed to double irradiation, total dosage 900 r (400 + 500 r). The second irradiation was conducted five and 15 days after the first.

The average life span of the mice in these groups was identical (273 ± 4 and 285 ± 14 minutes), which supports the abovecited observation that the conditions of irradiation (single or double) and time intervals between irradiations do not in-

resistance depends neither upon fractionation of the dose nor upon an increase in time intervals between irradiations. The following experiments were conducted to examine these problems.

In the second series of experiments the animals were subjected to double irradiation of the same total dosage — 750 r (400 + 350 r) — the time interval between exposures two and five days. In this short period of time about 60 and 75% of the reversible radiation injuries is repaired (cf. Diagram 1). The average life span of these mice after contamination by the toxin (339 ± 13 and 342 ± 13 minutes) essentially did not differ from the group exposed to single irradiation. Consequently, the degree of lesion of immunobiological resistance of the organism essentially does not depend upon the conditions of irradiation.

In the third series of experiments the influence of dosage and increased time interval between irradiations up to a period of time necessary for full reparation of reversible radiation alterations brought about by the first irradiation was examined. The animals of this series were exposed to double irradiation, total dosage 900 r (400 + 500 r). The second irradiation was conducted five and 15 days after the first.

The average life span of the mice in these groups was identical (273 ± 4 and 286 ± 14 minutes), which supports the abovecited observation that the conditions of irradiation (single or double) and time intervals between irradiations do not in-

fluence the degree of lesion of the organism's immunobiological state. However, the average life span of the animals in the third series exposed to 900 r (279 ± 4.9 minutes) was considerably less than that of the animals in the second series exposed to 750 r (355 ± 14 minutes). This statistically significant difference ($R = 0.99$) in the life span after contamination by a toxin is apparently linked with a difference in the doses of preliminary irradiation. With an increase of the irradiation dose the degree of lesion of immunobiological resistance of the organism grew, which also led to a greater decrease in the life span.

Our experimental material made it possible to establish that in an irradiated organism and for a period of time after restoration of radioresistance and reparation of reversible radiation injuries, lesions remain in the mechanism of immunobiological resistance for an exceedingly protracted and still unestablished period of time. This mechanism determines the ability of the organism to counteract the influence of bacterial toxins. Manifestation of these lesions, which was displayed in the reduction of the average life span after contamination by the toxin, *Vas. perfringens*, increased when the irradiation dose was increased. However, the conditions of the influence of radiation (single or double) and increasing the time intervals between irradiations

did not influence the degree of reduction of the organism's immunity. This allows us to associate lesions in the mechanism of immunobiological resistance with the irreversible component of radiation sickness.

It was shown earlier (A. V. Popov with co-authors) that for a protracted time after irradiation of the animals simultaneously with the lowering of immunobiological reactivity, increased hemopoiesis hidden by a normal or anemic pattern of peripheral blood was noted as well as an increase in thyroid activity of the thyroid gland, which was not accompanied by growth of the organ. Continued introduction of thyroid led to normalization of the state of immunobiological resistance and hemopoiesis and brought the histological structure of the thyroid gland and thyroglobulin formation closer to that of healthy animals. Probably, manifestations of the irreversible component of radiation sickness are specifically linked with the state of the endocrine system. Hormone therapy decreases these manifestations and possibly accelerates partial or complete restoration of these radiation injuries, which are repaired with the greatest of difficulty.

Bibliography

- 1) S.N. Aleksandrov, K.F. Galkovskaya, Meditainskaya radiologiya (Medical Radiology), 1959, No 11, page 15.
- 2) I. F. Kovalyov, Meditainskaya radiologiya (Medical Radiology), 1956, No 4, page 7.
- 3) V. I. Korogodin, G. G. Polikarpov, ibid., 1958, No 4, page 79.
- 4) M. A. Lagun, Tezisy dokl. na vtoroi mezhinstitutской konferentsii po voprosam radiatsionnoi mikrobiologii i immunologii (Theses and Reports at the Second Interinstitute Conference on Problems of Radiation Microbiology and Immunology), Moscow, 1960, page 46.
- 5) E. Blair, Materialy mezhdunarodnoi konferentsii po mirnomu ispol'zovaniyu atomnoi energii, (Material of the international conference on peaceful utilization of atomic energy), Moscow, 1958, T 11, page 147.
- 6) G. Davidson, Biologicheskiye posledstviya obshchego gamma-oblucheniya cheloveka (Biological Consequences of General Gamma-irradiation of Man), Moscow, 1960.
- 7) H. Kohn, R. Kallman, Radiation Resistance, 1957, vol 6, page 137; 329.
- 8) S. Rothermel, K. Woodard, J. Storer, ibid., 1956, vol 5, page 433.
- 9) J. Spalding, V. Strang, F. Worman, ibid., 1960, vol 13, page 415.
- 10) S. Storer, P. Harris, J. Furchner, et al, ibid., 1957, vol 6, page 188.
- 11) S. Tyler, S. Stearner, ibid., 1957, vol 7, page 253; 1960, vol 12, page 301.